Volume 13 , Number S-1 January, 2002

NSV Public NSWHEALTH Better Health Good Health Care Health Bulletin Supplement State Health Publication PH 010206

Trends in the prescribing of stimulant medication for the treatment of Attention Deficit Hyperactivity Disorder in children and adolescents in NSW

NSW DEPARTMENT OF HEALTH

Copyright © NSW Department of Health, January 2002

This work is copyright. It may be reproduced in whole or in part, subject to the inclusion of an acknowledgment of the source and no commercial usage or sale.

State Health Publication No: PH010206

ISBN 0734733690

suggested citation:

Salmelainen P. Trends in the prescribing of stimulant medication for the treatment of Attention Deficit Hyperactivity Disorder in children and adolescents in NSW. Sydney: NSW Department of Health, 2002.

produced by:

Pharmaceutical Services Branch Public Health Division NSW Department of Health PO Box 103 Gladesville NSW 1675 Australia

Tel: 61 2 9879 3214 Fax: 61 2 9859 5165

further copies of this publication can be obtained by contacting:

Better Health Centre Publication Warehouse Locked Bag 5003 Gladesville NSW 2111 Australia

Tel: 61 2 9816 0452 Fax: 61 2 9816 0492

or from the Public Health Division Web site at www.health.nsw.gov.au





Table of Contents

	EXECUTIVE	SUMMA	RY	4
ı.	INTRODUC	TION		6
2.	BACKGROU	JND		8
	2.1	Diagnosis,	symptoms and causes of ADHD	8
	2.1.1	_	and symptoms	
	2.1.2	_	ADHD	
	2.2	Prevalence	e of ADHD	10
	2.3	Treatment	s for ADHD	12
	2.3.1	Psychosoc	ial intervention	12
	2.3.2	Pharmaco	therapy	13
	2.4	Restriction	ns on stimulant prescribing in NSW	17
	2.4.1		pproval	
	2.4.2	Individual	patient approval	17
	2.5	Aim of thi	s study	19
3	METHODS			20
٥.	3.1		Trends in the number of children and adolescents treated	20
	3.1	section 1:	with stimulant medication for the first time	20
	3.2	Section 2	Trends in the rate of children and adolescents treated	20
	3.2	Section 2.	with stimulant medication	20
	3.3	Section 3:	Characteristics of children and adolescents treated with stimulant medication	21
	3.4	Section 4:	Two- to three-year olds treated with stimulant medication	22
	3.5	Section 5:	Attrition from stimulant treatment	22
	3.6	Caveats		23
4.	RESULTS	••••		24
	4.1	Section 1:	Trends in the number of children and adolescents	
			treated with stimulant medication for the first time	24
	4.1.1	Overall tr	end	24
			Number of children with ADHD commenced on	
			stimulant medication for the first time by year of	
			commencement, I 990 to 2000	22
	4.1.2	Age at cor	mmencement	24
		Figure 2:		
			stimulant medication for the first time, by age and year at	
		F: -	commencement, 1990 to 2000	25
		Figure 3:	Percent of children with ADHD commenced on	
			stimulant medication for the first time, by age and year at commencement, 1990 to 2000	24
	4.1.3	Sev	Commencement, 1990 to 2000	
	т.1.5		Number of children with ADHD commenced on	20
		1 1841 C 1.	stimulant medication for the first time by sex, 1990 to	

		2000	27
4.1.4	Drug		26
	Figure 5:	Number of children with ADHD commenced on	
		stimulant medication for the first time, by drug and year	
		at commencement, 1990 to 2000	27
4.2	Section 2:	Trends in the rate of children and adolescents treated	
		with stimulant medication	
4.2.1			28
	Figure 6:	Rate per 1,000 NSW resident population of children	
		treated with stimulant medication for ADHD by sex and	
	_	year, 30 June, 1990 to 2000	
4.2.2	J		28
	Figure 7:	Rate per 1,000 NSW resident population of children	
		treated with stimulant medication for ADHD by age and	
	_	year, 30 June, 1990 to 2000	
4.2.3	-		28
	Figure 8:	Rate per 1,000 NSW resident population of children	
		treated with stimulant medication for ADHD by drug and	20
4.5		year, 30 June, 1990 to 2000	30
4.3	Section 3:	Characteristics of children and adolescents treated with	21
421	A 1.0	stimulant medication	
4.3.1	_	ex	31
	Table 1:	Number and percentage of children treated with	
		stimulant medication for ADHD as at 1 December 2000 by age and sex	21
	Table 2:	, -	3 1
	rable 2:	Rate per 1,000 NSW resident population of children treated with stimulant medication for ADHD as at	
		I December 2000 by age and sex	32
	Figure 9:	Rate per 1,000 NSW resident population of children	52
	rigure 7.	treated with stimulant medication for ADHD as at	
		I December 2000 by single year of age and sex	32
4.3.2	Age and d	rug	
	•	Per cent of children treated with stimulant medication	
		for ADHD as at 1 December 2000 by age and drug	33
4.3.3	Age and d	ose	
	-	Average daily dose (in tablets) for children treated with	
		stimulant medication for ADHD as at 1 December 2000	
		by age and drug	34
	Figure 12:	Per cent of children treated with stimulant medication	
	J	for ADHD as at I December 2000 by age and daily dose	34
4.3.4	Health are	ea of residence	35
		Rate per 1,000 NSW resident population of children	
	0	treated with stimulant medication for ADHD as at	
		I December 2000 by health area of residence and sex	36
	Figure 14:	Rate per 1,000 NSW resident population of children	
	-	aged 5-6 years treated with stimulant medication for	
		ADHD as at I December 2000 by health area of	
		residence and sex	36
	Figure 15:	Rate per 1,000 NSW resident population of children	
		aged 7-11 years treated with stimulant medication for	

			ADHD as at 1 December 2000 by health area of residence and sex	37
		Figure 16:	Rate per 1,000 NSW resident population of children aged 12–15 years treated with stimulant medication for ADHD as at 1 December 2000 by health area of residence and sex	
		Figure 17:	Rate per 1,000 NSW resident population of children aged 16–17 years treated with stimulant medication for ADHD as at 1 December 2000 by health area of residence and sex	
	4.3.5	Duration	of stimulant treatment	
		Table 3:	Number and percentage of children treated with stimulant medication for ADHD as at 1 December 2000 by age and duration of continuous treatment	
	4.4	Section 4:	Two- to three-year olds treated with stimulant medication	
		Table 4:	Characteristics of children aged 2–3 years Initial authorities issued for stimulant medication for ADHD, I January 1999 to 30 June 2000	
		Case I:	'Michael', aged 2 years and 8 months	
		Case 1:	'Phillip', aged 2 years and 8 months	
		Case 3:	'Kylie', aged 2 years and 7 months	
		Case 4:	'Harry', aged 3 years and 7 months	
		Case 5:	'Daniel', aged 3 years and 8 months	
		Case 6:	'Peter', aged 3 years and I month	
	4.5		Attrition from stimulant treatment	
		Figure 18:	last prescription	
5.	DISCUSSIO	N		47
6.	REFERENCI	ES		53
	6.1	Useful into	ernet resources	57
7 .	APPENDIX	•••••		58
8.	ACKNOWL	EDGEME	NTS	65